

Lattice QCD Calculations of the Sigma Commutator

D. B. Leinweber*, A. W. Thomas†, and S. V. Wright‡

*Department of Physics and Mathematical Physics
and Special Research Centre for the Subatomic Structure of Matter,
University of Adelaide, Adelaide 5005, Australia*

Abstract

As a direct source of information on chiral symmetry breaking within QCD, the sigma commutator is of considerable importance. With recent advances in the calculation of hadron masses within full QCD it is of interest to see whether the sigma commutator can be calculated directly from the dependence of the nucleon mass on the input quark mass. We show that provided the correct chiral behaviour of QCD is respected in the extrapolation to realistic quark masses one can indeed obtain a fairly reliable determination of the sigma commutator using present lattice data. Within two-flavour, dynamical-fermion QCD the value obtained lies in the range 45 to 55 MeV.

In the quest to understand hadron structure within QCD, small violations of fundamental symmetries play a vital role. The sigma commutator, σ_N :

$$\sigma_N = \frac{1}{3} \langle N | [Q_{i5}, [Q_{i5}, \mathcal{H}]] | N \rangle \quad (1)$$

(with Q_{i5} the two-flavour ($i=1, 2, 3$) axial charge) is an extremely important example. Because Q_{i5} commutes with the QCD Hamiltonian in the chiral SU(2) limit, the effect of the double commutator is to pick out the light quark mass term from $\mathcal{H}$:

$$\sigma_N = \langle N | (m_u \bar{u}u + m_d \bar{d}d) | N \rangle \quad (2)$$

Neglecting the very small effect of the $u-d$ mass difference we can write Eq. (2) in the form

$$\sigma_N = \langle N | \bar{m} (\bar{u}u + \bar{d}d) | N \rangle \quad (3)$$

$$= \bar{m} \frac{\partial M_N}{\partial \bar{m}} \quad (4)$$

*dleinweb@physics.adelaide.edu.au

†athomas@physics.adelaide.edu.au

‡swright@physics.adelaide.edu.au

with $\bar{m} = (m_u + m_d)/2$. Equation (4) follows from the Feynman-Hellman theorem [3].

While there is no direct experimental measurement of σ_N , the value inferred from world data has been 45 ± 8 MeV [4] for some time. Recently there has been considerable interest in this value because of progress in the determination of the pion-nucleon scattering lengths [5, 6] and new phase shift analyses [7, 8]. For an excellent summary of the sources of the proposed variations and the disagreements between various investigators we refer to the excellent review of Knecht [9]. For our purposes the experimental value is of limited interest as the full lattice QCD calculations upon which our work is based involve only two active flavours. Nevertheless, as a guide, the current work suggests that the best value of σ_N may be between 8 and 26 MeV larger than the value quoted above [9].

Numerous calculations of σ_N have been made within QCD motivated models [10] and there has been considerable work within the framework of chiral perturbation theory [11]. However, direct calculations of σ_N within QCD itself have proven to be difficult. Early attempts [12] to extract σ_N from the quark mass dependence of the nucleon mass in quenched QCD (using Eq.(4)) produced values in the range 15 to 25 MeV. Attention subsequently turned to determining σ_N by calculating the scalar matrix element of the nucleon $\langle N | \bar{u}u + \bar{d}d | N \rangle$. There it was discovered that the sea quark loops make a dominant contribution to σ_N [14, 15]. These works, based on quenched QCD simulation, found values in the 40 to 60 MeV range, which are more compatible with the experimental values quoted earlier.

On the other hand, the most recent estimate of σ_N , and the only one based on a two-flavour, dynamical-fermion lattice QCD calculation, comes from the SESAM collaboration. They obtain a value of 18 ± 5 MeV [13], through a direct calculation of the scalar matrix element $\langle N | \bar{u}u + \bar{d}d | N \rangle$. The discrepancy from the quenched results of Refs. [14, 15] is not so much an unquenching effect in the scalar matrix element but rather a significant suppression of the quark mass in going from quenched to full QCD. The difficulty in all approaches which evaluate $\langle N | \bar{u}u + \bar{d}d | N \rangle$ is that neither it nor $\bar{m}$ is renormalization group invariant. One must reconstruct the scale invariant result from the product of the scale dependent matrix element and the scale dependent quark masses. The latter are extremely difficult to determine precisely and are the chief source of uncertainty in this approach.

An additional difficulty in extracting σ_N from lattice studies is the need to extrapolate from quite large pion masses, typically above 500 or 600 MeV. An important innovation adopted by Dong *et al.*, but not by the SESAM collaboration, was to extrapolate the computed values of $\langle N | \bar{u}u + \bar{d}d | N \rangle$ using a form motivated by chiral symmetry, namely $a + b\bar{m}^{\frac{1}{2}}$. On the other hand, the value of b used was *not* constrained by chiral symmetry and higher order terms of the chiral expansion were not considered. Furthermore, since the work was based on a quenched calculation, the chiral behaviour implicit in the lattice results involves incorrect chiral coefficients [16].

Our work is motivated by recent, dramatic improvements in computing power which, together with the development of improved actions [17], mean that we now have accurate calculations of the mass of the nucleon within *full QCD* (for two flavours) as a function of $\bar{m}$ down to $m_\pi \sim 500$ MeV. (Since m_π^2 is proportional to $\bar{m}$ over the range studied we choose to display all results as a function of m_π^2 .) In addition, CP-PACS has recently published a result at $m_\pi \sim 300$ MeV, albeit with somewhat large errors. Provided that one has control over the extrapolation of this lattice data to the physical pion mass, $m_\pi \equiv \mu = 140$ MeV, one can calculate σ_N by evaluating Eq. (4) at the physical pion mass. Note that this approach has the important advantage over the calculation of the scalar density that one *only* needs to work with renormalization group invariant quantities. We therefore turn to a consideration of the method of extrapolation.

The lattice data for the nucleon mass calculated by UKQCD [1] and CP-PACS [2] is shown in Fig. 1. Both groups cite a 10% uncertainty in setting the lattice scale, so we have scaled the former down and the latter up by 5% so that the data sets are consistent. Over almost the entire range of m_π^2 , the data shows a dependence on quark mass that is essentially linear. However, the preliminary point at $m_\pi^2 \sim 0.1$ GeV² suggests some curvature in the low mass region. This is indeed expected on the basis of chiral symmetry with the leading non-analytic (LNA) correction (in terms of $\bar{m}$) being proportional to m_π^3 ($\bar{m}^{3/2}$):

$$\delta M_N^{\text{LNA}} = \gamma^{\text{LNA}} m_\pi^3, \quad \gamma^{\text{LNA}} = \frac{-3g_A^2}{32\pi f_\pi^2}. \quad (5)$$

These observations led the CP-PACS group to extrapolate their data with the simple, phenomenological form:

$$M_N = \tilde{\alpha} + \tilde{\beta} m_\pi^2 + \tilde{\gamma} m_\pi^3. \quad (6)$$

The corresponding fit to the combined data set, using Eq. (6), is shown as the short-dashed curve in Fig. 1 and the parameters $(\tilde{\alpha}, \tilde{\beta}, \tilde{\gamma}) = (0.912, 1.69, -0.761)$ (the units are appropriate powers of GeV). This yields a value for the sigma commutator, $\sigma_N^{(p)} = 29.7$ MeV, where the superscript stands for “phenomenological”.

The difficulty with this purely phenomenological analysis was discussed in Ref. [18]. That is, the value of $\tilde{\gamma} = -0.761$ is almost an order of magnitude smaller than the model independent LNA term, $\gamma^{\text{LNA}} = -5.60$ GeV⁻². Clearly this presents some concern when evaluating σ_N , because of the derivative required. An alternative approach to this problem was recently suggested by Leinweber et al. [18]. They realised that the pion loop diagrams, Fig. 2(a) and 2(b) not only yield the most important non-analytic structure in the expression for the nucleon mass, but amongst all the possible meson baryon states which contribute to the nucleon mass within QCD, they alone give rise to a significant variation of the nucleon mass as $m_\pi \rightarrow 0$. In Ref. [18] it was suggested that one should extrapolate M_N as a function of quark mass using:

$$M_N = \alpha + \beta m_\pi^2 + \sigma_{NN}(m_\pi, \Lambda) + \sigma_{N\Delta}(m_\pi, \Lambda), \quad (7)$$




Figure 1: Nucleon mass calculated by CP-PACS (solid points) and UKQCD (open points), as a function of m_π^2 , both are scaled by 5% to improve consistency. The solid curve is a fit to Eq. (7) with a 1.225 GeV dipole form factor, while the dashed curve is the same fit using a sharp cut-off form factor (θ -function). The short-dash curve is a fit to Eq. (6), and the long-dash curve is a fit to Eq. (7) *excluding* the lowest data point. The vertical line indicates the physical pion mass.

where σ_{NN} and $\sigma_{N\Delta}$ are the self-energy contributions of Figs. 2(a) and 2(b), respectively, using a sharp cut-off in momentum, $\theta(\Lambda - k)$. The full analytical expressions for σ_{NN} and $\sigma_{N\Delta}$ are given in Ref. [18]. For our purposes it suffices that they have precisely the correct LNA and next-to-leading non-analytic behaviour required by chiral perturbation theory as $m_\pi \rightarrow 0$. In addition, $\sigma_{N\Delta}$ contains the correct, square root branch point ($\sim [m_\pi^2 - (M_\Delta - M_N)^2]^{\frac{3}{2}}$) at the $\Delta - N$ threshold, which is essential for extrapolations from above the $\Delta - N$ threshold.

Fitting Eq. (7) to the data, including the point near 0.1 GeV², gives the dot-dash curve in Fig. 1 ($(\alpha, \beta, \Lambda) = (1.42, 0.564, 0.661)$). The corresponding value of σ_N is 54.6 MeV and the physical nucleon mass is 870 MeV. Omitting the lowest data point from the fit yields the long-dash curve in Fig. 1 ($(\alpha, \beta, \Lambda) = (1.76, 0.386, 0.789)$) with $\sigma_N = 65.8$ MeV. Clearly the curvature associated with the chiral corrections at low quark mass is extremely important in the evaluation of σ_N .

In order to estimate the error in the extracted value of σ_N we would need to have the full data set on a configuration by configuration basis. As this is not




Figure 2: One-loop pion induced self energy of the nucleon.

Scaling		σ_N		
CP-PACS	UKQCD	Dipole	Sharp	Cubic
5	-5	47.2 ± 1.8	54.6 ± 2.0	29.7
10	0	48.1 ± 1.9	54.9 ± 2.0	28.6
0	-10	45.4 ± 1.9	54.3 ± 1.9	31.0

Table 1: Sigma Commutator Values. The **Dipole** and **Sharp** results were calculated with our preferred form of $\alpha + \beta m_\pi^2 + \sigma_{NN}(\Lambda, m_\pi) + \sigma_{N\Delta}(\Lambda, m_\pi)$ with either a dipole form factor for the $N\pi$ vertex or a θ -function. The values of dipole parameter (Λ_D) were (1.225, 1.250, 1.175) GeV. The **Cubic** results are for the $\alpha + \beta m_\pi^2 + \gamma m_\pi^3$ extrapolation function, with γ *unconstrained* by chiral symmetry – as explained in the text this produces an unreliable value for σ_N .

available, the errors that we quote are naive estimates only. The extracted value of σ_N is very well determined by the present data, the result being 54.6 ± 2.0 MeV. Since the process of setting the physical mass scale via the string tension is thought to have a systematic error of 10%, one might naively expect this to apply to σ_N . However, *all* masses in the problem including the pion (or quark) mass, as well as that of the nucleon, scale with the lattice parameter a . It turns out that when one uses Eq. (4) at the physical pion mass (which means a slightly different value of $\bar{m}a$ if a changes), the value of σ_N is extremely stable. If, for example, one raises the CP-PACS data by 15% and the UKQCD data by 5% (instead of 5% and -5% , respectively) the value of σ_N shifts from 54.6 ± 2.0 to 55.2 ± 2.1 MeV. We present calculations in Table 1 that show, for a variety of scalings of the lattice data, how stable our results are.

The remaining issue, for the present data, is the model dependence associated with the choice of a sharp cut-off in the pionic self-energies. Our investigations in Ref. [18] showed that Eq. (7) could reproduce the dependence of M_N on m_π^2 within the cloudy bag model, and that it could also describe the dependence of pion self-energy terms calculated with dipole form factors. Thus we believe that any model satisfying the essential chiral constraints and fitting the lattice data should give essentially the same answer. We checked this by numerically fitting the lattice data (solid curve) with the form of Eq. (7) but with σ_{NN} and $\sigma_{N\Delta}$ calculated with dipole form factors of mass Λ_D at all pion-baryon vertices. Since the preferred phenomenological form of the $N\pi$ form factor is a dipole, we regard the dipole result shown in the first line of Table 1 as our best estimate, namely $\sigma_N = 47.2 \pm 1.8$ MeV with fit parameters $(\alpha, \beta, \Lambda_D) = (2.02, 0.398, 1.225)$. A

remaining source of error is that, although the lattice results were calculated with an improved action, there still is an error associated with the extrapolation to the infinite volume, continuum limit. The importance of the inclusion of the correct chiral behaviour is clearly seen by the fact that it increases the value of the sigma commutator from the 30 MeV of the unconstrained cubic fit to around 50 MeV.

Clearly an enormous amount of work remains to be done before we will fully understand the structure of the nucleon within QCD. It is vital that the rapid progress on improved actions and faster computers continue and that we have three flavour calculations within full QCD at masses as close as possible to the physical quark masses. Nevertheless, it is a remarkable result that the present lattice data for dynamical-fermion, two-flavour QCD, yields such a stable and accurate answer for the sigma commutator, an answer which is already within the range of the experimental values. The implications of this result for models of hadron structure need to be explored urgently.

One of us (SVW) would like to acknowledge helpful discussions with Tom Cohen at an early stage of this work. We would also like acknowledge helpful comments from Chris Allton, Craig Roberts and Robert Perry. This work was supported in part by the Australian Research Council.

References

- [1] **UKQCD** Collaboration, C. R. Allton *et. al.*, [hep-lat/9808016](#).
- [2] **CP-PACS** Collaboration, S. Aoki *et. al.*, *Phys. Rev.* **D60**, 114508 (1999), [hep-lat/9902018](#).
- [3] R. P. Feynman, *Acta Phys. Polon.* **24** (1963) 697–722;
R. P. Feynman, *Phys. Rev.* **56** (1939) 340;
H. Hellman, *Einführung in die Quantenchemie*. Deutick Verlag, Leipzig, 1937;
S. T. Epstein, *Amer. J. Phys.* **22** (1954) 614.
- [4] J. Gasser, H. Leutwyler and M. E. Sainio, *Phys. Lett.* **B253**, 252 (1991).
- [5] D. Sigg *et al.*, *Nucl. Phys.* **A609** (1996) 269;
D. Chatellard *et al.*, *Nucl. Phys.* **A625**, 855 (1997).
- [6] T. E. Ericson, B. Loiseau and A. W. Thomas, “Precision determination of the πN scattering lengths and the charged πNN coupling constant”, [hep-ph/9907433](#).
- [7] M. M. Pavan, R. A. Arndt, I. I. Strakovsky and R. L. Workman, “New result for the pion nucleon sigma term from an updated VPI/GW πN partial-wave and dispersion relation analysis”, [nucl-th/9912034](#).
- [8] D. V. Bugg, “Summary of the Conference”, to be published in the proceedings of 8th International Symposium on Meson-Nucleon Physics

and the Structure of the Nucleon (MENU 99), Zuoz, Switzerland, 16-20 Aug 1999.

- [9] M. Knecht, [hep-ph/9912443](#).
- [10] I. Jameson, A. A. Rawlinson and A. W. Thomas, *Austral. J. Phys.* **47**, 45 (1994);
R. E. Stuckey and M. C. Birse, *J. Phys. G* **G23**, 29 (1997),
[hep-ph/9602312](#).
- [11] U. Meissner, “Chiral dynamics: Status and perspectives”,
[hep-ph/9810276](#);
V. Bernard, N. Kaiser and U. G. Meissner, *Z. Phys.* **C60**, 111 (1993),
[hep-ph/9303311](#).
- [12] See for example, S. Cabasino *et al.*, *Nucl. Phys. B* (Proc. Suppl.) **20**, 399 (1991).
- [13] **SESAM** Collaboration, S. Gusken *et al.*, *Phys. Rev.* **D59**, 054504 (1999),
[hep-lat/9809066](#).
- [14] S. J. Dong, J. F. Lagae and K. F. Liu, *Phys. Rev.* **D54**, 5496 (1996),
[hep-ph/9602259](#).
- [15] M. Fukugita, Y. Kuramashi, M. Okawa and A. Ukawa, *Phys. Rev.* **D51**, 5319 (1995), [hep-lat/9408002](#).
- [16] S.R. Sharpe, *Phys. Rev.* **D56**, 7052 (1997), [hep-lat/9707018](#);
J.N. Labrenz and S.R. Sharpe, *Phys. Rev.* **D54**, 4595 (1996),
[hep-lat/9605034](#).
- [17] B. Sheikholeslami and R. Wohlert, *Nucl. Phys.* **B259** (1985) 609;
M. Alford, W. Dimm, P. Lepage, *Phys. Lett.* **B361**, 87 (1995);
P. Hasenfratz, *Nucl. Phys.* (Proc. Suppl.) **B63**, 53 (1998).
- [18] D. B. Leinweber, A. W. Thomas, K. Tsushima, and S. V. Wright, “Baryon masses from lattice QCD: Beyond the perturbative chiral regime”,
[hep-lat/9906027](#). To appear in *Phys. Rev.* **D**.